Cosmology and the origin of structure

Rocky I: The universe observed

Rocky II: The growth of cosmological structure


Rocky III: Inflation and the origin of perturbations

Rocky IV: Dark matter and dark energy


Academic Training Lectures

Rocky Kolb


Fermilab, University of Chicago, & CERN




Missing Pieces




Dark Matter






$$\frac{v^2}{R} = \frac{GM_{\text{GALAXY}}}{R^2}$$


measure v and $R \Longrightarrow M_{\text{GALAXY}}$





Rotation curves

CO – central regions Optical – disks HI – outer disk & halo











Gravitational Lens Galaxy Cluster 0024+1654

Hubble Space Telescope • WFPC2





The evolved spectrum



Most of the universe is dark!


Modified Newtonian dynamics


Planets


Mass disadvantaged stars


brown red white


Black holes















Microlensing black-hole candidates




Run 1339 Col 4 Field 75

SteHodesainst Reachbold les In the Solar Neighborhood (with Jim Chisholm & Scott Dodelson)


Model spectral energy distributions:

$$L = \int \frac{dv}{v} v L_v$$
 $v = \begin{cases} v = v \\ v = v \end{cases}$ (contribution per decade)




Sloan: a 2.5m telescope & instruments to

- 1. image the sky to 23rd mag in 5 colors (10⁹ objects)
- 2. take the spectra of 10^6 objects (mostly extragalactic)








SDSS

2.5 m telescope


data acquisition system



SDSS camera



Color-color space




Color-color cut + RASS detection

SDSS Early Data Release = 3.7 million objects

SDSS color cut = 150,000






ROSAT

Röntgen Satellite X-Ray Observatory Germany/US/UK 1990-1999

 $0.1 \text{ keV} \le E \le 2.4 \text{ keV}$

ROSAT effective area



Color-color cut + RASS detection


SDSS Early Data Release = 3.7 million objects

SDSS color cut = 150,000

SDSS + RASS = 47

- 47 is a manageable number (can examine each individually)
- 7 targeted for spectroscopy by SDSS (5 stars + 2 QSOs)
- Can define measure of how far from stellar locus in 4 color-color spaces





Most of the universe is @@IK!

- Modified Newtonian dynamics
- Planets
- Mass disadvantaged stars
 brown red white
- Black holes
- The weight of space

gravitational microlensing

Most of the universe is dark!

- Modified Newtonian dynamics
- Planets
- Mass disadvantaged stars
 brown red white
- Black holes
- The weight of space
- Fossil remnant of the big bang

gravitational microlensing

Neutrinos?

- Neutrinos are known to exist three active + sterile?
- Neutrinos are strongly suspected to have mass

LSND O(eV) ATM.: $O(10^{-2} eV)$ SOLAR: $O(10^{-3} eV)$


Massive neutrinos contribute to the mass density

$$\Omega_{\nu\bar{\nu}}h^2 = \frac{m_{\nu}}{90 \text{ eV}}$$


Ballard Firehouse

January 26 February 9
\$3.00

The evolved spectrum



Projected SDSS BRG



NEUTRINO MASS AND MIXING IMPLIED BY UNDERGROUND DEFICIT OF LOW ENERGY MUON-NEUTRINO EVENTS

John G. LEARNED, Sandip PAKVASA, and Thomas J. WEILER 1

Department of Physics and Astronomy, University of Hawaii at Manoa, Honolulu, HI 96822, USA

Received 14 March 1988

Recent observations of a deficit of cosmic ray muon-neutrino interactions in underground detectors suggest that the muon neutrinos may have oscillated to another state. We examine possible neutrino mass and mixing patterns, and their implications for vacuum and matter effects on solar neutrinos, on neutrinos passing through the earth, and on terrastrial neutrino beams. By invoking the see-saw mechanism of neutrino mass generation, we draw inferences on closure of the universe with neutrino masses, on the number of generations, on t-quark and fourth generation masses, and on the Peccei-Quinn symmetry breaking scale. Testable predictions are suggested.

(d) Relic tau neutrinos have sufficient energy density to close the university

(6)

flux is delo estimate icreased by

nly,

(7)


otating the pasis. Since ng angle is

(8)

und (from are forced

and muon-neutrinos (not antineutrinos) coming through the earth at $E_v \sim 50 - 150$ GeV have matterenhanced oscillations and the muon-neutrinos down/up flux ratio should be even larger than the nonmatter-enhanced expectation (for energies ~1 TeV and larger oscillations are suppressed in the earth). Detectors capable of distinguishing μ^- from μ^+ have increased sensitivity to matter effects [10]. (c) A 40% v₁₁ ↔ v₂ conversion should be observable in a detector at a distance L from the neutrino source for the integrated flux satisfying $E/\text{GeV} \lesssim 0.04 L/\text{km}$. (d) Relic tau neutrinos have sufficient energy density to close the university [11] (thus favoring hot dark matter over cold); the tau-neutrino mass may be determined from the time spread of events from a galactic supernova. (e) There are only three generations: the mass of a fourth-generation heavy lepton is bounded from below by the UA1 data [12] and from

Cold thermal relics





Cold thermal relics


$$\Omega_X h^2 \sim \langle \sigma_A v \rangle^{-1}$$


$$\sigma_{\scriptscriptstyle A} \Leftrightarrow \Omega_{\scriptscriptstyle X}$$

$$\sigma_{\scriptscriptstyle A} \Leftrightarrow \sigma_{\scriptscriptstyle S}$$










CMSSM


 m_0 = universal scalar mass $m_{1/2}$ = gaugino mass $\tan \beta$ = ratio of vevs $|\mu|$ determined

Ellis, Olive, Santoso



 $m_{1/2}$

Ellis, Olive, Santoso



Thermal WIMP: interaction & mass limit


 Ω_X depends on the anihilation strength $(\Omega_X \propto \sigma_A^{-1})$ $\Omega_X \approx 1$ annihilation strength \approx electroweak scale annihilation strength \rightarrow interaction strength

$$\sigma_A \le \frac{8\pi}{m_V^2}$$
 unitarity limit to the cross section

$$\Omega_X \le 1 \Longrightarrow M_X \le 200 \,\mathrm{TeV}$$

Thermal WIMP: Interaction strength determined Mass undetermined (but < 200 TeV)

Nonthermal relics



Expanding universe particle creation



(Arnowit, Birrell, Bunch, Davies, Deser, Ford, Fulling, Grib, Hu, Kofman, Mostepanenko, Page, Parker, Starobinski, Unruh, Vilenkin, Wald, Zel'dovich,...)

first

density perturbations from inflation gravitational waves from inflation

(Guth & Pi; Starobinski; Bardeen, Steinhardt, & Turner; Hawking; Rubakov; Fabbi & Pollack; Allen)


new application: dark matter

(Chung, Kolb, & Riotto; Kuzmin & Tkachev)

- require (super)massive particle "X"
- stable (or at least long lived)
- initial inflationary era followed by radiation/matter

WIMPZILLA production

Chung, Kolb, Riotto (also Kuzmin & Tkachev)



 $\Omega_X \approx 1$ for $M_X/M_{\text{INFLATON}} \approx 1 \Rightarrow M_X \approx 10^{10}$ to 10^{15}GeV


Superheavy particles

Inflaton mass (in principle measurable from gravitational wave background, guess $10^{12}~{\rm GeV}$) may signal a new mass scale in nature.

Other particles may exist with mass comparable to the inflaton mass.

Superheavy relic (wimpzilla) characteristics:

- Supermassive: $10^9 10^{19}$ GeV (~ 10^{12} GeV?)
- abundance may depend only on mass
- abundance may be independent of interactions
 - sterile?
 - electrically charged?
 - strong interactions?
 - weak interactions?
- unstable (lifetime > age of the universe)?



Decay: Ultra High Energy Cosmic Rays

Annihilate: Galactic Center, Sun

Isocurvature Perturbations: Structure Formation, CMB


Direct Detection: Bulk, Underground Searches

Dark Matter

WIMP

or WIMPZILLA





Particle Dark Matter Candidates

- neutrinos (hot dark matter)
 sterile neutrinos, gravitinos (warm dark matter)
 LSP (neutralino, sneutrino, ...) (cold dark matter)
 axion, axion clusters
 WIMPZILLA
- solitons (B-balls; Q-balls; Odd-balls,....)

 - •

- Origin of structure:
 a complex natural phenonenon
- Gravitational instability of perturbations from inflation:
- a simple, elegant, compelling explanation

"For every complex natural phenomenon there is a simple, elegant, compelling, wrong explanation."

- Tommy Gold

What We "Know" *

The matter density is dominated by cold dark matter, which we know nothing about!

The perturbations arise from inflationary dynamics, which depends on particle physics at high energies, which we know nothing about!

The universe is dominated by a cosmological term (dark energy, funny energy, quintessence, polenta, cosmological constant, cosmoillogical constant,), which we know less than nothing about!

^{*}It ain't what you don't know, it's what you know that ain't so!

Cosmology and the origin of structure

Rocky I: The observed universe

Rocky II: The growth of cosmological structure

Rocky III: Inflation and the origin of perturbations

Rocky IV: Dark matter and dark energy

Academic Training Lectures

Rocky Kolb

Fermilab, University of Chicago, & CERN